

AMG™ Coupling Kit, 200 nm Magnetic Particles

AI-LNPCKMP-7.2

This AMG™ Coupling Kit contains particles that are fully activated with Mix&Go™ and are ready to use, with buffer solutions known to work for a majority of applications. Protocols should be optimised to meet individual requirements.

Materials Supplied	
Cat. No.	Component
A-CMPLNMP	Activated 200 nm Magnetic Particles
A-CMPCBC1	Coupling Buffer
A-CMPBBB3	Blocking Buffer
A-CMPSBA1	Storage Buffer
n/a	1.5 mL microcentrifuge tubes

Additional Materials Required
Protein of interest
Pipettes
Magnetic separator for tubes (capable to separate 200 nm magnetic particles)
Tube rotator
Vortex mixer
Microcentrifuge for 1.5 mL tubes
Sonication bath with fresh deionised water (minimum power 60W)

Specifications	
Ordering Information	A-LNPCKMP-10 (10 Reactions) A-LNPCKMP-30 (30 Reactions)
Storage	Store at 2°C - 8°C. Remaining materials should be retained in the supplied container and sealed for future use.
Stability	Particles coupled by the user should be assessed for individual use and storage stability conditions, as this can vary depending on the protein and conditions used.
Applications	Particles are compatible with the majority of existing uses and protocols. This allows for easy substitution of the AMG™ Coupling Kit into your application of choice.
Supplied Surface	Particle: 200 nm, Superparamagnetic, 70% Magnetite. Concentration: 10 mg/mL (1% w/v solids).
Additives	Product contains ProClin® 300 as preservative.
Regulatory	For laboratory use only.

Compatibility	
Buffers	It is recommended to use the included buffers as the presence of certain materials will interfere with and reduce the coupling efficiency and storage stability. Mix&Go products are not compatible with high concentrations of Phosphates. For coupling, it is important to dilute the phosphate concentration, with the provided Coupling Buffer, to ≤ 2.5 mM as per the Chemical Compatibility Table on page 4.
Tubes	It is recommended to use low binding polypropylene microcentrifuge tubes.
Temperature	Do not freeze or expose to temperatures exceeding 60°C. Room temperature is defined as 20°C - 25°C.
Protein Concentration	Protein coupling concentration is best optimised as this can vary depending on the protein used. The recommended concentration range for coupling antibody is 25 - 100 µg/mg particles, with a determined maximal binding capacity up to 100 µg/mg particles.
Magnet	When separating magnetic particles from supernatant, it is recommended to put the tube containing the particles on the magnetic separator until the supernatant is clear (up to 5 minutes), and carefully remove the supernatant so as not to disturb the particle pellet. Low strength magnets may take longer to form a pellet or may leave residue on the tube.

Safety
Standard safety precautions exercised when handling laboratory reagents should be adhered to. Refer to the product MSDS for safety precautions.

Other Products
For more information on Anteo products please visit www.anteotech.com .

Warnings/Hazards
Product is not guaranteed DNase, RNase or endotoxin free. The user must determine the suitability of the product for specific uses.

AMG™ Coupling Kit, 200 nm Magnetic Particles

AI-LNPCKMP-7.2

Procedure		Time
Coupling Preparation		
1.	Allow all components to come to room temperature	
2.	Vortex particles	10 seconds
3.	Sonicate particles	5 minutes
4.	Aliquot 100 µL (1 mg) of Activated 200 nm Magnetic Particles (Cat No. A-CMPLNMP) to a provided 1.5 mL tube	
Washing with Coupling Buffer		
5.	Pulse in a microcentrifuge to ensure particles are at the bottom of the tube before separation	
6.	Place the tube on the magnetic separator to separate particles from solution	5 minutes
7.	Remove and discard the supernatant from the tube and then remove the tube from the magnet	
8.	Resuspend the particles in 100 µL of Coupling Buffer (Cat. No. A-CMPCBC1)	
9.	Vortex particles	10 seconds
ⓘ	Particles may aggregate and appear to stick on the tube during coupling. This may be part of the normal process whereby the surface during coupling is interacting with your protein of interest. Particles can be dispersed by sonication in a bath sonicator.	
10.	Repeat washing steps for a total of 2 washes	
Coupling		
11.	Prepare 100 µL of antibody at the required final concentration in Coupling Buffer (Cat. No. A-CMPCBC1) in a fresh tube (provided)	
12.	Pulse the particle tube in a microcentrifuge to ensure particles are at the bottom of the tube before aspiration	
13.	Aspirate all of the prepared particles	
14.	Add the particles to the antibody solution	
15.	Vortex particles	10 seconds
16.	Sonicate particles	5 minutes
ⓘ	Total volume is 200 µL total volume and particles are at 5 mg/mL.	
17.	Incubate at room temperature on a tube rotator to keep the particles in suspension	60 minutes
Blocking		
18.	Vortex particles	10 seconds
19.	Add 20 µL of the Blocking Buffer (Cat. No. A-CMPBBB3) to the tube	
20.	Vortex particles	10 seconds
21.	Incubate at room temperature on a tube rotator to keep the particles in suspension	60 minutes
Washing with Storage Buffer		
22.	Vortex particles	10 seconds
23.	Pulse in a microcentrifuge to ensure particles are at the bottom of the tube before separation	
24.	Place the tube on the magnetic separator to separate particles from solution	5 minutes
25.	Remove and discard the supernatant from the tube and then remove the tube from the magnet	
26.	Resuspend the particles in 200 µL of Storage Buffer (Cat. No. A-CMPSBA1)	
27.	Vortex particles	10 seconds
28.	Repeat washing steps for a total of 2 washes	
Storage		
29.	Pulse in a microcentrifuge to ensure particles are at the bottom of the tube before separation	
30.	Place the tube on the magnetic separator to separate particles from solution	5 minutes
31.	Remove and discard the supernatant from the tube and then remove the tube from the magnet	
32.	Resuspend the particles in 100 µL of Storage Buffer (Cat. No. A-CMPSBA1)	
33.	Vortex particles	10 seconds
34.	Sonicate particles	5 minutes
35.	Vortex particles	10 seconds
36.	Sonicate particles	5 minutes
ⓘ	Particles may be used immediately or stored at 2°C - 8°C. Final particle concentration is 10 mg/mL. Resuspend particles (step 33 - 36) prior to use.	

AMG™ Coupling Kit, 200 nm Magnetic Particles

AI-LNPCKMP-7.2

Procedure							
Coupling Preparation							
Washing with Coupling Buffer							
Coupling							
Blocking							
Washing with Storage Buffer							
Storage							

AMG™ Coupling Kit, 200 nm Magnetic Particles

AI-LNPCKMP-7.2

Chemical Compatibility Table			
	Coupling	Storage	Assay
EDTA	X	X	≤ 10 mM
Phosphates (PBS 10 mM Phosphates, 150 mM NaCl)	≤ 2.5 mM	X	≤ 20 mM
Tween20®	≤ 0.25%	X	≤ 0.5%
DMSO	≤ 15%	X	≤ 30%
pH	Use Coupling Buffer (pH 5.2) provided	Use Storage Buffer provided	ND
Urea	≤ 2 M	X	≤ 2 M
Saline	≤ 150 mM	≤ 150 mM	≤ 150 mM
Sodium Azide	≤ 0.1%	≤ 0.1%	≤ 0.1%
ProClin 300®	≤ 0.1%	≤ 0.1%	≤ 0.1%
Other Proteins	Co-coupling	X	OK

Final concentration of chemicals during coupling.

Contact		
Website: www.anteotech.com	Anteo Technologies Pty Ltd	Office: +61 7 3219 0085
Technical Support: support@anteotech.com	Unit 4, 26 Brandl Street	Fax: +61 7 3219 0553
Bulk Pricing & Sales Enquiries: sales@anteotech.com	Eight Mile Plains QLD 4113	
	Australia	

All Anteo Products are sold as general purpose reagents for general laboratory and research uses only. Anteo Products are not intended for diagnostic and/or therapeutic purposes and no Anteo Product may be administered to humans. Anteo does not make any representation or warranty that Anteo Products comply with all laws and regulations that may be applicable to Customer's use of any Anteo Product.

Anteo warrants that Anteo Products will conform to the specifications set forth on the applicable Anteo Product Certificate of Analysis ("**Anteo's Limited Warranty**"). The Certificate of Analysis may be obtained by contacting Anteo. Anteo's Limited Warranty is wholly conditioned on the proper use of Anteo Products in the applications for which they are intended, and Anteo makes no warranty (express, implied or statutory) for Anteo Products that are modified; subjected to accident, misuse, neglect, unauthorized repair or improper tampering, testing or storage; and/or used or handled contrary to Anteo's instructions as set forth on the Anteo Product label and/or in this insert.

Unless otherwise expressly provided in this document or in these Terms and Conditions of Sale, Anteo disclaims all warranties, conditions and guarantees, whether written, express, implied, statutory or otherwise, including but not limited to, the implied warranties or guarantees of merchantability and fitness for particular purpose.

ProClin® 300 is a registered trademark of Rohn and Haas Company.

Mix&Go™ is a registered Australian trademark of Anteo Technologies Pty Ltd.